INDEX

Africa, produce buying in, 280–8.
Andrews, F. W., control of nut grass in the Sudan Gezira, 214–22.

Barley, malting, cultivation of, in England, 65–79; grading of samples, 65–7, 67–9; influence of: lodging, 77–8; manuring, 70–2; previous crop, 74–6; season, 67; soil, 69–70; time of sowing, 72–4; variety of seed, 70; weather at harvest, 76–7.
Beckley, V. A., compost-making in Kenya, 311–18.
Bidner, N., *see* Ravikovith, S.
Bisschop, J. H. R., bionomic studies on indigenous and exogenous cattle in the semi-arid regions of the Union of S. Africa, 138–47.
Brenchley, W. E., the weed problem in non-rotational wheat-growing, 126–37.

Campbell, J. T., *see* Ward, A. H.
Cattle, beef, records of performance for, 259–63; methods of assessing market grade, 260–2; suggested procedure, 262–3.
— breeding of, in Tanganyika Territory, 11–22; crossing of Zebu cows with imported bulls: Afrikander, 14; Ayrshire, 15–16; Friesian, 14–15; Krishna Valley, 14; Shorthorn, 14; degeneration of imported herds in tropics, 12–13, 14 sqq.; indigenous herds, 11–13; influence of environment, 12, 13, 16; of heredity, 13, 19–20; of temperature, 17–19; selection and breeding from indigenous stock, 13, 20–1.
— dairy sires, evaluation of, in New Zealand, 249–58; herd-recording, 249–50; methods of compiling and issuing sire-surveys, 250–1; results and analysis of 'final' sire-surveys, 252, 253–5; 'sire indices', 251–2, 253, 255.
— indigenous and exogenous, in semi-arid regions of South Africa, 138–47; bionomic experiments at Armoedsvlakte, 138–9; conditions at Armoedsvlakte, 139–411; macro-morphological research methods, 141–5; micro-morphological methods, 145–7.
— influence of atmospheric temperature on haemoglobin, &c., of blood, 97–100; blood-changes in Indian Nellore, Holstein-Friesian, and their F1 crosses, in Philippines, 98–9; non-adaptability of imported stock to high temperatures and high humidity, 99.
— method of assaying genetic differences in adaptability to tropical and subtropical climates, 190–8; experimental animals and conditions, 191–2; N-concentration in urine, 194–5; rate of water-vaporization from skin, 194; rectal temperatures, 195–7; respiratory rate, 193; water in faeces, 195.
Ceylon, experiments on tea in, 269–79.
Christidis, B. G., viability of cotton seed as affected by its moisture-content, 148–58.
Clover, effects of soil salinity on, in Yezreel Valley, 207–14.
— in Kikuyu-grass pasture, Kenya, 102 sqq.
Compost-making in Kenya, 311–18; Adco process, 311–12; cereal residues, 311, 312, 314–15; coffee-cherry pulp, 311, 315–16; coffee parchment, 317; composting *in situ*, 317; Indore process, 313–14, 317; large-scale practices, 314–17; Napier grass, 314, 315–16; sisal waste, 316–17; town wastes, 317; water-supply and wetting, 312, 313–14, 315; weeds, 313, 314.
Cotton industry, control of, in Uganda, 281–4.
Cotton seed, viability of, as affected by moisture-content, 148–58; effect of drying-temperature, 149, 154; of storage after drying, 150, 155, 157; field experiments, 155, 156–7; germinability, 149, 151 sqq.; rate of moisture-loss, 151, 155.
Crowther, E. M., *see* Haines, W. B.

Davies, Wm., winter forage in Britain: possible use of Pampas and Tussac grasses, 289–94.
De Vries, O., determination of the potash requirement of soils, 295–300.
Deighton, T., ventilation, temperature and humidity in piggeries and other farm buildings, 319–22.
Design, modern experimental, and its function in plant selection, 223–30; advantage of using many varieties in field trials, 223; classification of designs, 224; lattice designs, 224–9.
Diseases, plant, due to minor-element deficiencies, 85–96, 199–206. *Boron*: brown heart (raan) of swedes, turnips, 89–90; brown spotting of apricots, 92; cracked stem of celery, 91; heart rot of celery, 91, and of sugar-beet, 89; internal cork of apples, 91; *topzieckte* of tobacco, 89, 90–1. *Copper*: die-back (exanthema) in citrus, 199, 200; reclamation diseases, 199, 200–1. Diagnosis of deficiency diseases, 86–8. *Manganese*: 92–5; grey speck of oats, 92–8; marsh spot of peas, 87, 94–5; Pahala

XUM

blight of sugar-cane, 93; speckled yellows of mangold and sugar-beet, 93. *Zinc*: bronzing of tung-oil trees, 204–5; little leaf (rosetting) of fruit trees, 202; mottle leaf of citrus, 204; white bud of maize, 202–3.
Donald, H. P., breeding policy in relation to performance-testing of pigs, 185–9.

Eden, T., studies in the yield of tea. IV. Effect of cultivation and weeds on crop growth, 269–79.
Edwards, D. C., reaction of Kikuyu-grass herbage to management, 101–10.

Falcon, Prudencio R., *see* Manresa, M.
Farming, primitive (Mumuye), in N. Nigeria, 51–5; break-up of tribal life, 55; crops and stock, 52–3; cultivations and harvesting, 54; farming guilds, 53; land tenure, 51–2; manuring, 53; soil erosion, 53; tribal cults, 53–4.
Feeding-stuffs, Britain's supplies of, 231–48; allocation of, to live stock, 240–3; changes in live-stock population, 235, 236; choice of supplies in war-time, 237–40; food-requirements of live stock, 232, 245; imported animal products, 247, 248; ploughing-up campaign, 239–40; pre-War imports of feeding-stuffs, 232–7, 245, 246, 247; pre-War supplies, 231–2, 244.
Finney, D. J., the Little Hoos Field experiment on the residual values of certain manures, 111–25.
Forage crops, new, for Palestine, 1–5; cow-pea, 3; field-pea, 2; horse-bean, 2; Jerusalem artichoke, 4–5; *Lathyrus ochrus*, 2; *Pencillarea*, 4; pumpkin, 4; Sudan grass, 5; sunflower, 4; sweet lupin, 2–3; teff grass, 3–4; *Vicia fulgens*, 3; 'wonder forage', 4.
Forage, winter, in Britain, 289–95; methods of conserving summer-grown fodder, 289–90; possible use of pampas grass, 290–2; of tussac grass, 292–3.
Fraser, A. H. H., *see* Robertson, D.
French, M. H., cattle-breeding in Tanganyika Territory and some developmental problems relating thereto, 11–22.

Garner, H. V., cultivation of malting barley in England, 65–79.
Gomez, Francisco, *see* Manresa, M.
Grass, methods of conserving, 288–9; 'foggaging', 288–9; *in situ*, 290; silage, 288.
— products, shortage of supplies in winter in Britain, 232.
Ground-nut industry in N. Nigeria, economic aspects of, 39–50; exports, 39, 40; markets and marketing, 44–6; prices, 42–3; producing areas, 39–40; production and factors affecting, 40–4; State aids, 49–50; transport, 44, 46–9.

Haines, W. B., manuring *Hevea*, Pt. III, results on young buddings in British Malaya, 169–84.
Hammond, John, *see* McMeekan, C. P.
Herd-recording in New Zealand, 349–50.
Hurvitz, S., new forage crops for Palestine, 1–5.

India, varieties of potato grown in, 80–4.

Kenya, compost-making in, 311–18.
— Kikuyu-grass pasture in, 101–10.
Kikuyu-grass pasture, reaction of, to management, 101–10; composition after grazing, 104; after regular cutting, 105–6; before grazing, 103; experimental procedure, 101–3.

Lambs, influence of a moderate worm-burden on the growth and health of, 323–5; experimental details, 323; symptoms and results of infestation, 324–5.
Liversage, V., control of produce buying in Africa, 280–8.

Ma, Ruh-hwa, and Kao, Lih-ming, a factorial experiment on rice culture, 23–33.
McMeekan, C. P., relation of environmental conditions to breeding and selection for commercial types in pigs, 6–10.
Malaya, manurial experiments on *Hevea*, 169–84.
Manresa, Miguel, influence of atmospheric temperature upon haemoglobin and other constituents of the blood of cattle, 97–100.
Manures, experiment on residual values of, 111–25; experimental details, 111–12; manuring, 113–15; nitrogen in crops, 123–4; phosphorus in crops, 124–5; yields on control plots, 115–17, on manured plots, 117–23.
Mauritius, sugar-cane varieties in, 301–10.
Meat animals, records of performance for, 259–68; beef cattle, 259–63; pigs, 266–7; sheep, 263–5; utilization of recorded data, 267–8.
Minor elements, deficiencies of, in agricultural and horticultural crops, 85–96, 199–206; boron, 88–92; copper, 199–202; manganese, 92–5; molybdenum, 205; zinc, 202–5.
Moorland, reclamation experiments in Scotland, 56–64; early work, 56–7; nature and

composition of *Scirpus* and *Eriophorum* peats, 59–60; recent work in Lanark, 58–60, in Lewis, 59–60; reclamation costs, 63, cropping, 62–3, cultivations, 61–2, drainage, 60–1, liming, 61, manuring, 61.

New South Wales, wool-production in, 34–8.
New Zealand, evaluation of dairy sires in, 249–58.
Nichols, J. E., some aspects of selection of Merino stock for wool-production under pastoral conditions, 34–8.
Nigeria, Northern, ground-nut industry of, 39–50.
— — primitive farming in, 51–5.
Nut grass, control of, in the Sudan Gezira, 215–22; climate of the Gezira, 215; habitat, root-system, &c., of nut grass, 215, 216–17; treatment and results, 218–21.

Ogg, W. G., the reclamation of moorland, Pt. II, reclamation experiments in Scotland, 56–64.

Pal, B. P., varieties of potatoes grown in India, 80–4.
Palestine, new forage crops for, 1–5.
—, Yezreel Valley, soil salinity in, 207–14.
Pampas grass, 290–2.
Pease, M. S., present position of poultry research, 326–8.
Peat-land in Scotland, reclamation of, 56–64.
Peats, composition of moorland, 59.
Performance records for meat animals, 259–68.
Philippines, non-adaptability of imported pure-bred cattle in, 97–100.
Piggeries and other farm buildings, microclimate of, 319–22; computation of heat losses and gains, 319–20; humidity, 321; temperature, 320–1, 322; ventilation, 321–2.
Pig husbandry, co-ordinated experiments in, 159–67; data recorded, 160; experimental programme, 166–7; layout of experiments, 161; results, analysis of, 161–6.
Pigs, breeding policy in relation to performance-testing, 185–9; pedigrees of Large Whites in breeders' sale catalogues, 185; proportion of home-bred to not home-bred, 186–8.
— records of performance for, 266–7; suggested procedure, 266–7.
— relation of environmental conditions to breeding and selection for commercial types of, 6–10; bacon and lard types, 6 sqq.; effects of differential feeding, 7, 9; experimental method, 7, 8; importance of genetic selection, 9.
Piper, C. S., symptoms and diagnosis of minor-element deficiencies in agricultural and horticultural crops. Pt. I, 85–96; Pt. II, 199–206.
Plant selection, *see* Design, modern experimental.
Potash requirement of soils, comparison of methods of determining, 295–300; experimental data, 295–6; results, 296–300.
Potato varieties grown in India, 80–4.
Poultry, present position of research in, 326–8; disease and mortality, 327; influence of genetical research, 327–8; of statistical research, 326; organization of research in general, 326–7; research in Germany, 326.
Produce buying in Africa, 280–8; advantages of control, 280–1; is monopoly control desirable? 280; marginal costs, 284–6; monopoly control in Uganda, 281–4; official price-fixing, 286–8.

Ravikovitch, S., effects of soil salinity on clover in the Yezreel Valley, 207–14.
Reyes, N. C., *see* Manresa, M.
Rhoad, A. O., a method of assaying genetic differences in the adaptability of cattle to tropical and subtropical climates, 190–8.
Rice, a factorial experiment on, 23–33; design and layout, 23–4, 25; empty panicles, 25, 27–9; interactions, 24, 27, 30–1; yield effects of spacing, 22, 28, 29; time of heading, 25, 29–31; transplanting age, 24, 25 sqq.; varieties, 24, 26 sqq.
Robertson, D., influence of a moderate worm burden on the growth and health of lambs, 323–5.
Robertson, Ian M., *see* Ogg, W. G.
Rubber (*Hevea*), results of manuring young buddings of, 169–84; experimental technique, 169–72; fertilizer effects on growth of immature trees, 172–6, 177, on mature trees, 176, 177–8; replanting experiment, 182; responses to various phosphates, 181–3; yield effects, 178 sqq.
Russell, E. R., primitive farming in Nigeria: the Mumuye tribe, 51–5.

Sheep, records of performance for, 263–5; Christgau's method, 263–4; suggested procedure, 264–5.
— selection of Merino, for wool-production, 34–8.
Shorruck, R. W., co-ordinated experiments in pig husbandry, 159–67.
Soil cultivations; among Mumuye tribe, 54.
— — effect of, on weed populations, 127–34.
— — for control of nut grass, 217–21.

XUM

Soil cultivations; on reclaimed moorland, 61–2.
— — on tea soil, 269–77.
Soils, potash requirement of, 295–300.
— saline clover, in Yezreel Valley, Palestine, 207–14; chlorides and total salts in, 208–9; constituents of clover grown on, 209–10; ditto. on soil in Jordan valley, 210–11; movement of salts due to irrigation, 211–14.
Stevenson, G. C., sugar-cane varieties in Mauritius, 301–10.
Sudan (Anglo-Egyptian), control of nut grass in the Gezira, 215–22.
Sugar-cane, varieties of, in Mauritius, 301–10; early history, 303–4; Cheribon cane, 305–8; interspecific hybrid canes, 309; noble canes, 302 sqq.; Otaheite cane, 304–5; Penang (Salangore) cane, 308–9; present breeding trends, 310; Tanna cane, 308.

Tanganyika Territory, cattle-breeding in, 11–22.
Tea, effect of cultivation and weeds on crop growth, 269–79; effect of bulk manure on yields, 277–8; of clean weeding on soil conservation, 278–9; effects of cultivation and weeds on crop yields, 276–7; on root-distribution, 274–6; on soil resistance, 271–4; scheme of investigation, 270–1.
Turner, R., some economic aspects of the ground-nut industry of Northern Nigeria, 39–50.
Tussac grass, 292–3.

Uganda, monopoly control of cotton industry in, 281–4.
Union of S. Africa, indigenous and exogenous cattle in, 138–47.

Van Itallie, Th. B., *see* O. de Vries.

Ward, A. H., evaluation of dairy sires in New Zealand, 249–58.
Weeds, in tea soil, 269 sqq.
— on Broadbalk continuous-wheat field, Rothamsted, 126–37; effects of cultivations, fallowing, 127–34, and manuring, 135–6; germination periods, 128; numbers of viable buried seeds, 129, 131; recolonization, 132–4.
— use of, in compost-making, 313, 314.
Weil, J. W., *see* Garner, H. V.
Winters, L. M., records of performance for meat animals, 259–68.
Wool-production, selecting Merino stock for, in Australia, 34–8; fleece-weights in New South Wales, 34–6; limitations to genetic improvement, 37; stud flocks in W. Australia, 36.
Wright, Norman C., Britain's supplies of feeding-stuffs, 231–48.

Yates, F., modern experimental design and its function in plant selection, 223–30.

Zebu, short-horned, use of, for cross-breeding in Tanganyika, 12 sqq.
Zialcita, Lorenzo P., *see* Manresa, M.

XUM